Express Mail Label No.: EV781048075US Attorney Docket No.: 27353-501 UTIL

(Formerly 0623.0860002/LBB/Y-W) Date of Deposit: February 1, 2006

Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

(Previously presented) A method of generating a protein array, the method comprising: 1.

(a) inserting a marker DNA sequence in frame immediately following a start codon of

each of a plurality of target DNA sequences or immediately preceding a stop codon of each of a

plurality of target DNA sequences or both, to form a plurality of modified DNA sequences which

encode a plurality of modified amino acid sequences each comprising a marker moiety;

(b) expressing the plurality of modified amino acid sequences from the plurality of

modified DNA sequences;

(c) bringing the plurality of modified amino acid sequences into contact with a solid

support wherein the marker moiety of the plurality of modified amino acid sequences is able to

attach to the solid support, thereby generating a protein array, and

(d) washing said solid support to remove unbound proteins.

(Previously presented) The method as claimed in claim 1 wherein the marker moiety is a 2.

peptide sequence selected from the group consisting of:

a histidine tag; (a)

a complete protein or protein domain; and (b)

a maltose binding protein domain. (c)

3. (Previously presented) The method as claimed in claim 1 wherein the marker moiety

allows for purification of the individual proteins in the array.

4. (Previously presented) The method of claim 1 wherein the marker DNA sequence is

inserted such that the start or stop codon for each of the proteins is replaced.

5. (Canceled) An array prepared by a method as defined in any one of claims 1 to 4.

Page 2 of 11

6. (Canceled) An array as claimed in claim 5 wherein the components of the array are immobilised, eg to a solid surface.

7. (Canceled) An array as claimed in claim 6 wherein the individual proteins are immobilised by means of the tag moiety.

8. (Withdrawn) A method of screening one or more compounds for biological activity which comprises:

- (a) bringing said one or more compounds into contact with the array made according to any one of claims 1 to 4; and
 - (b) measuring binding of the one or more compounds to the proteins in the array.
- 9. (Withdrawn) A method of screening one or more proteins for specific protein-protein interactions which comprises the step of bringing said one or more proteins into contact with an array made according to any one of claims 1 to 4, and measuring binding of the one or more specific proteins with the proteins of the array.
- 10. (Withdrawn) A method of screening one or more proteins for specific interactions with one or more nucleic acid probes which comprises the step of bringing said one or more nucleic acid probes into contact with an array made according to any one of claims 1 to 4, and measuring binding of the probes to the proteins in the array.
- 11. (Withdrawn) A method for the rapid screening of a test compound, test protein or test nucleic acid, the method comprising:
- (a) contacting the test compound, test protein or test nucleic acid with a spatially defined array produced according to any one of claims 1-4 comprising a plurality of array bound proteins, with each array bound protein being at a different position on a solid support, wherein the plurality of array bound proteins comprises a plurality of different proteins expressed in a single species; and

(b) detecting any interaction between the array bound proteins and the test compound, test protein or test nucleic acid.

- 12. (Withdrawn) A method of screening for molecules which recognize each protein in the array, the method comprising:
- (a) contacting the molecules with a spatially defined array comprising a plurality of array bound proteins produced according to any one of claims 1-4, with each array bound protein being at a different position on a solid support, wherein the plurality of array bound proteins comprises a plurality of different proteins expressed in a single species; and
 - (b) detecting any interaction between the array bound proteins and the molecules.
- 13. (Previously presented) A method of generating an antibody array which comprises
- (a) bringing a protein array, made according to any one of claims 1 to 4, into contact with an antibody library, such that one or more proteins in the protein array bind to at least one antibody in the antibody library;
 - (b) removing any unbound antibodies; and
 - (c) immobilisation of those antibodies bound to proteins in the protein array.
- 14. (Withdrawn) A method for the screening of protein function or abundance which comprises the step of bringing an antibody array as defined in claim 13 into contact with a mixture of one or more proteins.
- 15. (Canceled) A method as claimed in any one of claims 8 to 10 and 13 which also comprises the step of first providing the protein array using a method as defined in any one of claims 1 to 4.
- 16. (Previously presented) The method of claim 1 wherein the marker DNA sequence is inserted immediately preceding a stop codon of a target DNA sequence by:
- (a) digesting the target DNA sequence such that it has a 5' overhang wherein the stop codon is comprised in the first three nucleotides counting from the 3' side of the overhang;

(b) annealing the marker DNA sequence to the overhang wherein the marker DNA

sequence comprises a sequence complementary to the first four nucleotides of the overhang

counting from the 3' side;

(c) ligating the marker DNA sequence to the target DNA sequence.

17. (Previously presented) The method of claim 1 wherein the marker DNA sequence is

inserted immediately following a start codon of a target DNA sequence by;

(a) digesting the target DNA sequence such that it has a 5' overhang wherein the start

codon is comprised in the first three nucleotides counting from the 3' side of the overhang;

(b) annealing the marker DNA sequence to the overhang wherein the marker DNA

sequence comprises a sequence complementary to the first four nucleotides of the overhang

counting from the 3' side;

(c) ligating the marker DNA sequence to the target DNA sequence.

18. (Previously presented) The method of any one of claims 1 to 4 wherein the protein array

comprises serine proteases, kinases or p450 enzymes.

19. (Previously presented) The method of any one of claims 1 to 4 wherein said plurality of

modified amino acid sequences are modified human amino acid sequences.

20. (Previously presented) The method of claim 1 wherein the marker moiety is selected

from the group consisting of FLAG and Strep.

21. (Previously presented) The method of claim 1 or 2 wherein the marker moiety is post-

translationally modified.

22. (Previously presented) The method of claim 21 wherein the post-translational

modification comprises the addition of a biotin or a lipid molecule.

Page 5 of 11

23. (Previously presented) The method of claim 1 wherein said modified amino acid sequences are folded into the correct conformation.

- 24. (Previously presented) The method of claim 1 wherein said inserting step inserts a marker DNA sequence in frame immediately following a start codon of each plurality target DNA sequence and immediately preceding a stop codon of each of a plurality of target DNA sequences, to form a plurality of modified DNA sequences which encode a plurality of modified amino acid sequences each comprising two marker moieties.
- 25. (Withdrawn) A method of generating a proteomic array of proteins of unknown amino acid sequences comprising the steps of:
 - (a) providing a cDNA library as a plurality of target DNA sequences; and
- (b) generating a protein array using the method of any of claims 1 to 4 to produce a proteomic array of proteins of unknown amino acid sequence.
- 26. (Currently amended) The method of claim 12, wherein the molecules are antibodies. A method of screening for antibodies which recognize each protein in the array, the method comprising:
- (a) contacting the antibodies with a spatially defined array comprising a plurality of array bound proteins produced according to any one of claims 1-4, with each array bound protein being at a different position on a solid support, wherein the plurality of array bound proteins comprises a plurality of different proteins expressed in a single species; and
 - (b) detecting any interaction between the array bound proteins and the antibodies.
- 27. (New) The method of claim 1, wherein the marker moiety provides a high-affinity attachment to the solid support.